

## Comment on “Solidification of a Supercooled Liquid in a Narrow Channel”

In a recent paper [1], Sabouri-Ghomi, et al. report on simulations of solidification in a channel geometry using a phase-field method. They conducted two series of simulations, one with very small surface tension (relative to the channel width), and the other at larger surface tension. This system has been studied extensively both analytically and numerically [2, 3, 4, 5, 6, 7]. The authors do not relate to the more recent of these studies [4, 5, 6, 7] which in fact explain many of the results the authors claim as surprising and can correct some misinterpretations presented.

Let us address the large surface tension case first. The authors find a transition from a tip widening instability for low undercooling,  $\Delta \lesssim 0.5$ , to a stable finger for  $\Delta \gtrsim 0.5$ . The tip widening instability, as noted by [4] is a result of the lack of steady-state solutions for sufficiently low  $\Delta$ . This threshold in  $\Delta$  is a function of surface tension  $d_0$  and anisotropy  $\epsilon$ . Thus, for example, at  $d_0 = 0.01$ , (in units where the channel width  $L = 2$ ) there are no steady-state solutions below  $\Delta \approx 0.62$  for zero anisotropy and below  $\Delta \approx 0.60$  at  $\epsilon = 0.1$  [6]. (Note that we define  $\epsilon$  as the anisotropy of the surface *tension* and not of the surface *energy*, as in Ref. [1], which is a factor of 15 smaller for 4-fold anisotropy.) This threshold decreases with  $d_0$ , going to 0 as  $d_0 \rightarrow 0$  and also with anisotropy. Above this threshold, there exist a bilaterally symmetric finger, which is the channel analogue of the free dendrite. We have extended the calculation [6] of the threshold to the parameters of the large surface tension case of Ref. [1], namely (in our units)  $d_0 = 0.00563$  and  $\epsilon = 0.75$ . The results are presented in Fig. 1, where we show the Péclet number, (dimensionless velocity),  $p$ , as a function of  $\Delta$ . We see that the transition from the unstable Saffman-Taylor branch to the dendritic branch occurs at  $\Delta = 0.49$ , in good agreement with the onset of stable dendritic growth reported in Ref. [1]. The stable pattern seen above the threshold is thus nothing but the well-known channel analogue of the dendrite [4, 6], and not a new type of solution. For example, we show in Fig. 2 the dendrite for  $\Delta = 0.55$ , which compares well with the pattern observed by the authors at this undercooling (see their Fig. 4). Also, this discussion should make clear that, contrary to the authors’ speculations, there is nothing special happening at  $\Delta = 0.5$ , which is of course the distinguished value in the Saffman-Taylor problem. We should also point out that at large  $\Delta$ , there also exist nonsymmetric fingers [4, 5, 6, 7], which have been termed parity-breaking dendrites, or doublons. The authors appear to have imposed reflection symmetry about the midline of their channel, which of course precludes seeing such solutions.

For small  $d_0$ , the authors only find fingers which slow down in time, and widen correspondingly. The surprise here is that the authors did not see stable translating




FIG. 1: Péclet number vs. supersaturation  $\Delta$  for symmetric solutions in a channel at  $d_0 = 0.00563$  and  $\epsilon = 0.75$




FIG. 2: Dendrite for  $\Delta = 0.55$  for parameters as in Fig. 1

dendrites. However, these simulations were performed at very small  $\Delta$ , so it is possible that again the authors are below the threshold for dendrites. We in fact suspect that the effective  $d_0$  for this series of runs is in fact much larger than the nominal value  $d_0 = 1.6 \times 10^{-5}$  quoted, since it is extremely difficult to adequately resolve the interface over such small scales, even with adaptive gridding.

Efim A. Brener

Institut für Festkörperforschung, Forschungszentrum Jülich  
D-52425 Jülich, Germany

David A. Kessler

Dept. of Physics, Bar-Ilan Univ.  
Ramat-Gan, IL52900 Israel

- [1] M. Sabouri-Ghomi, N. Provatas and M. Grant, Phys. Rev. Lett. **86**, 5084 (2001).
- [2] D. A. Kessler, J. Koplik and H. Levine, Phys. Rev. A **34**, 4980 (1986).
- [3] E. A. Brener, M. Geilikman and D. E. Temkin, Sov. Phys. JETP **67**, 1002 (1988).
- [4] E. A. Brener, H. Müller-Krumbhaar, Y. Saito and D. E. Temkin, Phys. Rev. E **47**, 1151 (1993).
- [5] T. Ihle and H. Müller-Krumbhaar, Phys. Rev. Lett. **70**, 3083 (1993); Phys. Rev. E **49**, 2972 (1994).
- [6] R. Kupferman, D. A. Kessler and E. Ben-Jacob, Physica **A213**, 451 (1995).
- [7] M. Ben Amar and E. A. Brener, Phys. Rev. Lett. **75**, 561 (1995); Physica **D98**, 128 (1996).